arXiv:1212.0619v1 [math.CO] 4 Dec 2012

Spectra of subdivision-vertex and subdivision-edge joins of graphs

Xiaogang Liu and Zuhe Zhang
Department of Mathematics and Statistics
The University of Melbourne
Parkville, VIC 3010, Australia
`xiaogliu.yzhang@gmail.com, zhang.zuhe@gmail.com`

Abstract

The subdivision graph $\mathcal{S}(G)$ of a graph G is the graph obtained by inserting a new vertex into every edge of G. Let G_1 and G_2 be two vertex disjoint graphs. The *subdivision-vertex join* of G_1 and G_2, denoted by $G_1\dot{\vee}G_2$, is the graph obtained from $\mathcal{S}(G_1)$ and G_2 by joining every vertex of $V(G_1)$ with every vertex of $V(G_2)$. The *subdivision-edge join* of G_1 and G_2, denoted by $G_1\underline{\vee}G_2$, is the graph obtained from $\mathcal{S}(G_1)$ and G_2 by joining every vertex of $I(G_1)$ with every vertex of $V(G_2)$, where $I(G_1)$ is the set of inserted vertices of $\mathcal{S}(G_1)$. In this paper we determine the adjacency spectra, the Laplacian spectra and the signless Laplacian spectra of $G_1\dot{\vee}G_2$ (respectively, $G_1\underline{\vee}G_2$) for a regular graph G_1 and an arbitrary graph G_2, in terms of the corresponding spectra of G_1 and G_2. These results enable us to construct infinite pairs of cospectral graphs.

Keywords: Spectrum, Cospectral graphs, Subdivision-vertex join, Subdivision-edge join

AMS Subject Classification (2010): 05C25, 05C50

1 Introduction

Throughout this article, all graphs considered are simple and undirected. Let $G = (V(G), E(G))$ be a graph with vertex set $V(G) = \{v_1, v_2, \ldots, v_n\}$ and edge set $E(G)$. The *adjacency matrix* of G, denoted by $A(G) = (a_{ij})_{n\times n}$, is an $n \times n$ symmetric matrix such that $a_{ij} = 1$ if vertices v_i and v_j are adjacent and 0 otherwise. Let $d_i = d_G(v_i)$ be the degree of vertex v_i in G and $D(G) = \mathrm{diag}(d_1, d_2, \ldots, d_n)$ be the diagonal matrix of vertex degrees. The *Laplacian matrix* and *signless Laplcacian matrix* of G are defined as $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$, respectively. Given an $n \times n$ matrix M, denote by

$$\phi(M; x) = \det(xI_n - M),$$

or simply $\phi(M)$, the characteristic polynomial of M, where I_n is the identity matrix of size n. In particular, for a graph G, we called $\phi(A(G))$ (respectively, $\phi(L(G))$, $\phi(Q(G))$) the *adjacency* (respectively, *Laplacian, signless Laplacian*) *characteristic polynomial* of G and its roots the *adjacency* (respectively, *Laplacian*, *singless Laplacian*) eigenvalues of G. The adjacency eigenvalues of G, denoted by $\lambda_1(G) \geq \lambda_2(G) \geq \cdots \geq \lambda_n(G)$, are called the *A-spectrum* of G. Similarly, the eigenvalues of $L(G)$ and $Q(G)$, denoted by $0 = \mu_1(G) \leq \mu_2(G) \leq \cdots \leq \mu_n(G)$ and $\nu_1(G) \leq \nu_2(G) \leq \cdots \leq \nu_n(G)$ respectively, are called the *L-spectrum* and *Q-spectrum* of G accordingly. Two graphs are said to be *A-cospectral* (respectively, *L-cospectral, Q-cospectral*) if they have the same A-spectrum (respectively, *L-spectrum*, *Q-spectrum*). It is well known that graph spectra store a lot of structural information about a graph. See [2,4,5] and the references therein to know more.

Up till now, many graph operations such as the disjoint union, the Cartesian product, the Kronecker product, the corona, the edge corona and the neighborhood corona have been introduced, and their spectra are computed in [2–7,9,10], respectively. It is well known [5] that the *subdivision graph* $\mathcal{S}(G)$ of a graph G is the graph obtained by inserting a new vertex into every edge of G. We denote the set of such new vertices by $I(G)$. In [8], the following new graph operations based on subdivision graphs are introduced.

Definition 1.1. The *subdivision-vertex join* of two vertex disjoint graphs G_1 and G_2, denoted by $G_1\dot{\vee}G_2$, is the graph obtained from $\mathcal{S}(G_1)$ and G_2 by joining each vertex of $V(G_1)$ with every vertex of $V(G_2)$.

Definition 1.2. The *subdivision-edge join* of two vertex disjoint graphs G_1 and G_2, denoted by $G_1\underline{\vee}G_2$, is the graph obtained from $\mathcal{S}(G_1)$ and G_2 by joining each vertex of $I(G_1)$ with every vertex of $V(G_2)$.

In [8], the A-spectra of $G_1\dot{\vee}G_2$ (respectively, $G_1\underline{\vee}G_2$), when G_1 and G_2 are both regular graphs, were computed in terms of the A-spectra of G_1 and G_2. As an application, the author constructed infinite family of new integral graphs (Graphs with their spectra only integers). In this paper, we will determine the A-spectra of $G_1\dot{\vee}G_2$ (respectively, $G_1\underline{\vee}G_2$) for a regular graph G_1 and an *arbitrary* graph G_2 in terms of that of G_1 and G_2 (see Theorems 2.1 and 3.1); this generalises [8, Theorems 1.1 and 1.2]. We also determine the L-spectra and the Q-spectra of $G_1\dot{\vee}G_2$ (respectively, $G_1\underline{\vee}G_2$) for a regular graph G_1 and an arbitrary graph G_2 (see Theorems 2.5, 2.8, 3.5 and 3.8). As we will see in Corollaries 2.4, 2.6, 2.10, 3.4, 3.6 and 3.10, our results on the spectra of $G_1\dot{\vee}G_2$ and $G_1\underline{\vee}G_2$ enable us to construct infinite pairs of cospectral graphs.

2 Spectra of subdivision-vertex joins

In this section, we determine the spectra of subdivision-vertex join with the help of the *coronal* of a matrix. The *M-coronal* $\Gamma_M(x)$ of an $n\times n$ matrix M is defined [3,9] to be the sum of the entries of the matrix $(xI_n - M)^{-1}$, that is,

$$\Gamma_M(x) = \mathbf{1}_n^T(xI_n - M)^{-1}\mathbf{1}_n,$$

where $\mathbf{1}_n$ denotes the column vector of dimension n with all the entries equal one.

It is well known [3, Proposition 2] that, if M is an $n\times n$ matrix with each row sum equal to a constant t, then

$$\Gamma_M(x) = \frac{n}{x-t}. \tag{2.1}$$

In particular, since for any graph G_2 with n_2 vertices, each row sum of $L(G_2)$ is equal to 0, we have

$$\Gamma_{L(G_2)}(x) = n_2/x. \tag{2.2}$$

2.1 A-spectra of subdivision-vertex joins

Theorem 2.1. *Let G_1 be an r_1-regular graph on n_1 vertices and m_1 edges, and G_2 an arbitrary graph on n_2 vertices. Then*

$$\phi\left(A(G_1\dot{\vee}G_2);x\right)=\phi(A(G_2);x)\cdot x^{m_1-n_1}\cdot\Big(x^2-n_1x\Gamma_{A(G_2)}(x)-2r_1\Big)\cdot\prod_{i=2}^{n_1}\Big(x^2-r_1-\lambda_i(G_1)\Big).$$

Proof. Let R be the incidence matrix of G_1. Then, with a proper labeling of vertices, the adjacency matrix of $G_1\dot{\vee}G_2$ can be written as

$$A(G_1\dot{\vee}G_2)=\begin{bmatrix}0_{n_1\times n_1} & R & J_{n_1\times n_2}\\ R^T & 0_{m_1\times m_1} & 0_{m_1\times n_2}\\ J_{n_2\times n_1} & 0_{n_2\times m_1} & A(G_2)\end{bmatrix},$$

where $J_{s\times t}$ denotes the $s\times t$ matrix with all entries equal one, and $0_{s\times t}$ denotes the $s\times t$ matrix with all entries equal zero.

Thus the adjacency characteristic polynomial of $G_1\dot{\vee}G_2$ is given by

$$\begin{aligned}\phi\left(A(G_1\dot{\vee}G_2);x\right) &= \det\begin{bmatrix}xI_{n_1} & -R & -J_{n_1\times n_2}\\ -R^T & xI_{m_1} & 0_{m_1\times n_2}\\ -J_{n_2\times n_1} & 0_{n_2\times m_1} & xI_{n_2}-A(G_2)\end{bmatrix}\\ &= \det(xI_{n_2}-A(G_2))\cdot\det(S)\\ &= \phi(A(G_2))\cdot\det(S),\end{aligned}$$

where

$$\begin{aligned}S &= \begin{pmatrix}xI_{n_1} & -R\\ -R^T & xI_{m_1}\end{pmatrix}-\begin{pmatrix}J_{n_1\times n_2}\\ 0_{m_1\times n_2}\end{pmatrix}(xI_{n_2}-A(G_2))^{-1}\begin{pmatrix}J_{n_2\times n_1} & 0_{n_2\times m_1}\end{pmatrix}\\ &= \begin{pmatrix}xI_{n_1}-\Gamma_{A(G_2)}(x)J_{n_1\times n_1} & -R\\ -R^T & xI_{m_1}\end{pmatrix}\end{aligned}$$

is the Schur complement [11] of $xI_{n_2}-A(G_2)$. It is well known [5] that $RR^T=A(G_1)+r_1I_{n_1}$. Thus, by applying (2.1), the result follows from

$$\begin{aligned}\det(S) &= x^{m_1}\cdot\det\left(xI_{n_1}-\Gamma_{A(G_2)}(x)J_{n_1\times n_1}-\frac{1}{x}RR^T\right)\\ &= x^{m_1}\cdot\left\{\det\left(xI_{n_1}-\frac{1}{x}RR^T\right)-\Gamma_{A(G_2)}(x)\cdot\mathbf{1}_{n_1}^T\operatorname{adj}\left(xI_{n_1}-\frac{1}{x}RR^T\right)\mathbf{1}_{n_1}\right\}\\ &= x^{m_1}\cdot\left(1-\Gamma_{A(G_2)}(x)\cdot\mathbf{1}_{n_1}^T\left(xI_{n_1}-\frac{1}{x}RR^T\right)^{-1}\mathbf{1}_{n_1}\right)\cdot\det\left(xI_{n_1}-\frac{1}{x}RR^T\right)\\ &= x^{m_1}\cdot\Big(1-\Gamma_{A(G_2)}(x)\cdot\Gamma_{\frac{1}{x}RR^T}(x)\Big)\cdot\prod_{i=1}^{n_1}\left(x-\frac{r_1}{x}-\frac{1}{x}\lambda_i(G_1)\right)\\ &= x^{m_1-n_1}\cdot\Big(x^2-2r_1-n_1x\cdot\Gamma_{A(G_2)}(x)\Big)\cdot\prod_{i=2}^{n_1}\Big(x^2-r_1-\lambda_i(G_1)\Big),\end{aligned}$$

where adj(M) is the adjugate matrix of M. □

Theorem 2.1 enables us to compute the A-spectra of many subdivision-vertex joins. In general, if we can determine the $A(G_2)$-coronal $\Gamma_{A(G_2)}(x)$, then we are able to compute the A-spectrum of $G_1\dot{\vee}G_2$. The following corollaries aim to illustrate this method.

Corollary 2.2. [8, Theorem 1.1] *Let G_1 be an r_1-regular graphs on n_1 vertices and m_1 edges, and G_2 an r_2-regular graphs on n_2 vertices. Then the A-spectrum of $G_1\dot{\vee}G_2$ consists of:*

(a) $\lambda_i(G_2)$, *for* $i = 2, 3, \ldots, n_2$;

(b) 0, *repeated* $m_1 - n_1$ *times;*

(c) $\pm\sqrt{r_1 + \lambda_j(G_1)}$, *for* $j = 2, 3, \ldots, n_1$;

(d) *the three roots of the equation*

$$x^3 - r_2x^2 - (n_1n_2 + 2r_1)x + 2r_1r_2 = 0.$$

Proof. (b) and (c) can be obtained readily from Theorem 2.1. Since G_2 is r_2-regular with n_2 vertices, (2.1) implies that

$$\Gamma_{A(G_2)}(x) = \frac{n_2}{x - r_2}.$$

The only pole of $\Gamma_{A(G_2)}(x)$ is $x = r_2$, which is equal to ${}_1(G_2)$. Thus, by Theorem 2.1, for each $i = 2, 3, \ldots, n_2$, $\lambda_i(G_2)$ is an eigenvalue of $G_1\dot{\vee}G_2$. Moreover, three more eigenvalues are obtained by solving

$$x^2 - 2r_1 - n_1x \cdot \frac{n_2}{x - r_2} = 0,$$

and this yields the eigenvalues in (d). □

Denote by $K_{p,q}$ the complete bipartite graph with $p, q \geq 1$ vertices in the two parts of its bipartition. Note that the case when $p = q$ is also covered by Corollary 2.2.

Corollary 2.3. *Let G be an r-regular graph on n vertices and m edges with $m \geq n$, and let $p, q \geq 1$ be integers. Then the A-spectrum of $G\dot{\vee}K_{p,q}$ consists of:*

(a) 0, *repeated* $m - n + p + q - 2$ *times;*

(b) $\pm\sqrt{r + \lambda_i(G)}$, *for* $i = 2, 3, \ldots, n$;

(c) *the four roots of the equation*

$$x^4 - \big(pq + n(p + q) + 2r\big)x^2 - 2npqx + 2pqr = 0.$$

Proof. It is known [9] that the $A(K_{p,q})$-coronal of $K_{p,q}$ is given by

$$\Gamma_{A(K_{p,q})}(x) = \frac{(p + q)x + 2pq}{x^2 - pq}.$$

The A-spectrum of $K_{p,q}$ [2,5] consists of $\pm\sqrt{pq}$ with multiplicity one, and 0 with multiplicity $p + q - 2$. Since $\pm\sqrt{pq}$ are the poles of $\Gamma_{A(K_{p,q})}(x)$, the result follows from Theorem 2.1 immediately. □

Up till now, many infinite families of pairs of A-cospectral graphs are generated by using graph operations (for example, [1, 9]). Now, we can use the subdivision-vertex join to obtain many A-cospectral graphs, as stated in the following corollary of Theorem 2.1.

Corollary 2.4. (a) *If G_1 and G_2 are A-cospectral regular graphs, and H is any graph, then $G_1 \dot{\vee} H$ and $G_2 \dot{\vee} H$ are A-cospectral.*

(b) *If G is a regular graph, and H_1 and H_2 are A-cospectral graphs with $\Gamma_{A(H_1)}(x) = \Gamma_{A(H_2)}(x)$, then $G \dot{\vee} H_1$ and $G \dot{\vee} H_2$ are A-cospectral.*

2.2 L-spectra of subdivision-vertex joins

Theorem 2.5. *Let G_1 be an r_1-regular graph on n_1 vertices and m_1 edges, and G_2 an arbitrary graph on n_2 vertices. Then*

$$\begin{aligned}\phi\left(L(G_1 \dot{\vee} G_2); x\right) &= x \cdot (x-2)^{m_1 - n_1} \cdot \Big(x^2 - (2 + r_1 + n_1 + n_2)x + 2n_1 + 2n_2 + n_1 r_1\Big) \\ &\quad \cdot \prod_{i=2}^{n_2} \Big(x - n_1 - \mu_i(G_2)\Big) \cdot \prod_{i=2}^{n_1} \Big(x^2 - \big(2 + r_1 + n_2\big)x + 2n_2 + \mu_i(G_1)\Big).\end{aligned}$$

Proof. Let R be the incidence matrix of G_1. Then the Laplacian matrix of $G_1 \dot{\vee} G_2$ can be written as

$$L(G_1 \dot{\vee} G_2) = \begin{bmatrix} (r_1 + n_2) I_{n_1} & -R & -J_{n_1 \times n_2} \\ -R^T & 2I_{m_1} & 0_{m_1 \times n_2} \\ -J_{n_2 \times n_1} & 0_{n_2 \times m_1} & n_1 I_{n_2} + L(G_2) \end{bmatrix}.$$

Thus the Laplacian characteristic polynomial of $G_1 \dot{\vee} G_2$ is given by

$$\begin{aligned}\phi\left(L(G_1 \dot{\vee} G_2); x\right) &= \det \begin{bmatrix} (x - r_1 - n_2) I_{n_1} & R & J_{n_1 \times n_2} \\ R^T & (x-2) I_{m_1} & 0_{m_1 \times n_2} \\ J_{n_2 \times n_1} & 0_{n_2 \times m_1} & (x - n_1) I_{n_2} - L(G_2) \end{bmatrix} \\ &= \det\big((x - n_1) I_{n_2} - L(G_2)\big) \cdot \det(S) \\ &= \det(S) \cdot \prod_{i=1}^{n_2} \Big(x - n_1 - \mu_i(G_2)\Big),\end{aligned}$$

where

$$S = \begin{pmatrix} (x - r_1 - n_2) I_{n_1} - \Gamma_{L(G_2)}(x - n_1) J_{n_1 \times n_1} & -R \\ -R^T & (x-2) I_{m_1} \end{pmatrix}$$

is the Schur complement [11] of $(x - n_1) I_{n_2} - L(G_2)$. Since G_1 is r_1-regular, we have $\lambda_i(G_1) =$

$r_1 - \mu_i(G_1)$, $i = 1, 2, \ldots, n_1$. Then

$$\begin{aligned}
\det(S) &= (x-2)^{m_1} \cdot \det\left((x - r_1 - n_2)I_{n_1} - \Gamma_{L(G_2)}(x-n_1)J_{n_1 \times n_1} - \frac{1}{x-2}RR^T\right) \\
&= (x-2)^{m_1} \cdot \left\{\det\left((x - r_1 - n_2)I_{n_1} - \frac{1}{x-2}RR^T\right)\right. \\
&\quad \left. -\Gamma_{L(G_2)}(x-n_1) \cdot \mathbf{1}_{n_1}^T \operatorname{adj}\left((x - r_1 - n_2)I_{n_1} - \frac{1}{x-2}RR^T\right)\mathbf{1}_{n_1}\right\} \\
&= (x-2)^{m_1} \cdot \det\left((x - r_1 - n_2)I_{n_1} - \frac{1}{x-2}RR^T\right) \\
&\quad \cdot \left(1 - \Gamma_{L(G_2)}(x-n_1) \cdot \mathbf{1}_{n_1}^T \left((x - r_1 - n_2)I_{n_1} - \frac{1}{x-2}RR^T\right)^{-1} \mathbf{1}_{n_1}\right) \\
&= (x-2)^{m_1 - n_1} \cdot \prod_{i=1}^{n_1} \left(x^2 - \big(2 + r_1 + n_2\big)x + 2n_2 + \mu_i(G_1)\right) \\
&\quad \cdot \left(1 - \Gamma_{L(G_2)}(x-n_1) \cdot \Gamma_{\frac{1}{x-2}RR^T}(x - r_1 - n_2)\right).
\end{aligned}$$

Hence, the result follows by applying (2.1) and (2.2) to $\det(S)$. □

Theorem 2.5 implies the following result.

Corollary 2.6. (a) *If G_1 and G_2 are L-cospectral regular graphs, and H is an arbitrary graph, then $G_1 \dot{\vee} H$ and $G_2 \dot{\vee} H$ are L-cospectral.*

(b) *If G is a regular graph, and H_1 and H_2 are L-cospectral graphs, then $G \dot{\vee} H_1$ and $G \dot{\vee} H_2$ are L-cospectral.*

(c) *If G_1 and G_2 are L-cospectral regular graphs, and H_1 and H_2 are L-cospectral graphs, then $G_1 \dot{\vee} H_1$ and $G_2 \dot{\vee} H_2$ are L-cospectral.*

Let $t(G)$ denote the number of spanning trees of G. It is well known [4] that if G is a connected graph on n vertices with Laplacian spectrum $0 = \mu_1(G) < \mu_2(G) \leq \cdots \leq \mu_n(G)$, then

$$t(G) = \frac{\mu_2(G) \cdots \mu_n(G)}{n}.$$

By Theorem 2.5, we can readily obtain the following result.

Corollary 2.7. *Let G_1 be an r_1-regular graph on n_1 vertices and m_1 edges, and G_2 an arbitrary graph on n_2 vertices. Then*

$$t(G_1 \dot{\vee} G_2) = \frac{2^{m_1 - n_1} \cdot (2n_1 + 2n_2 + n_1 r_1) \cdot \prod_{i=2}^{n_2}\left(n_1 + \mu_i(G_2)\right) \cdot \prod_{i=2}^{n_1}\left(2n_2 + \mu_i(G_1)\right)}{m_1 + n_1 + n_2}.$$

2.3 Q-spectra of subdivision-vertex joins

Theorem 2.8. *Let G_1 be an r_1-regular graph on n_1 vertices and m_1 edges, and G_2 an arbitrary graph on n_2 vertices. Then*

$$\begin{aligned}\phi\left(Q(G_1\dot{\vee}G_2);x\right) &= (x-2)^{m_1-n_1}\cdot\Big(x^2-(2+r_1+n_2)x+2n_2-n_1(x-2)\cdot\Gamma_{Q(G_2)}(x-n_1)\Big)\\ &\quad\cdot\prod_{i=1}^{n_2}\Big(x-n_1-\nu_i(G_2)\Big)\cdot\prod_{i=1}^{n_1-1}\Big(x^2-(2+r_1+n_2)x+2(r_1+n_2)-\nu_i(G_1)\Big).\end{aligned}$$

Proof. Let R be the incidence matrix of G_1. Then the signless Laplacian matrix of $G_1\dot{\vee}G_2$ can be written as

$$Q(G_1\dot{\vee}G_2)=\begin{bmatrix}(r_1+n_2)I_{n_1} & R & J_{n_1\times n_2}\\ R^T & 2I_{m_1} & 0_{m_1\times n_2}\\ J_{n_2\times n_1} & 0_{n_2\times m_1} & n_1I_{n_2}+Q(G_2)\end{bmatrix}.$$

The result follows by applying $RR^T=Q(G_1)$ and refining the arguments used to prove Theorem 2.5. □

Again, by applying (2.1), Theorem 2.8 implies the following result.

Corollary 2.9. *Let G_1 be an r_1-regular graphs on n_1 vertices and m_1 edges, and G_2 an r_2-regular graphs on n_2 vertices. Then*

$$\begin{aligned}\phi\left(Q(G_1\dot{\vee}G_2);x\right) &= (x-2)^{m_1-n_1}\cdot(x^3-ax^2+bx-4r_2n_2)\cdot\prod_{i=1}^{n_2-1}\Big(x-n_1-\nu_i(G_2)\Big)\\ &\quad\cdot\prod_{i=1}^{n_1-1}\Big(x^2-(2+r_1+n_2)x+2(r_1+n_2)-\nu_i(G_1)\Big),\end{aligned}$$

where $a=2+2r_2+r_1+n_1+n_2$ and $b=2n_1+2n_2+n_1r_1+2r_1r_2+2r_2n_2+4r_2$.

Theorem 2.8 can enable us to construct infinite Q-cospectral graphs.

Corollary 2.10. (a) *If G_1 and G_2 are Q-cospectral regular graphs, and H is a regular graph, then $G_1\dot{\vee}H$ and $G_2\dot{\vee}H$ are Q-cospectral.*

(b) *If G is a regular graph, and H_1 and H_2 are Q-cospectral graphs with $\Gamma_{Q(H_1)}(x)=\Gamma_{Q(H_2)}(x)$, then $G\dot{\vee}H_1$ and $G\dot{\vee}H_2$ are Q-cospectral.*

3 Spectra of subdivision-edge joins

In this section, we determine the spectra of subdivision-edge join.

3.1 A-spectra of subdivision-edge joins

Theorem 3.1. *Let G_1 be an r_1-regular graph on n_1 vertices and m_1 edges, and G_2 an arbitrary graph on n_2 vertices. Then*

$$\phi\left(A(G_1\underline{\vee}G_2);x\right)=\phi(A(G_2);x)\cdot x^{m_1-n_1}\cdot\Big(x^2-m_1x\Gamma_{A(G_2)}(x)-2r_1\Big)\cdot\prod_{i=2}^{n_1}\Big(x^2-r_1-\lambda_i(G_1)\Big).$$

Proof. Let R be the incidence matrix of G_1. Then the adjacency matrix of $G_1\underline{\vee}G_2$ can be written as

$$A(G_1\underline{\vee}G_2)=\begin{bmatrix}0_{n_1\times n_1} & R & 0_{n_1\times n_2}\\ R^T & 0_{m_1\times m_1} & J_{m_1\times n_2}\\ 0_{n_2\times n_1} & J_{n_2\times m_1} & A(G_2)\end{bmatrix}.$$

Thus the adjacency characteristic polynomial of $G_1\underline{\vee}G_2$ is given by

$$\begin{aligned}\phi\left(A(G_1\underline{\vee}G_2);x\right) &= \det\begin{bmatrix}xI_{n_1} & -R & 0_{n_1\times n_2}\\ -R^T & xI_{m_1} & -J_{m_1\times n_2}\\ 0_{n_2\times n_1} & -J_{n_2\times m_1} & xI_{n_2}-A(G_2)\end{bmatrix}\\ &= \det(xI_{n_2}-A(G_2))\cdot\det(S)\\ &= \phi(A(G_2))\cdot\det(S),\end{aligned}$$

where

$$S=\begin{pmatrix}xI_{n_1} & -R\\ -R^T & xI_{m_1}-\Gamma_{A(G_2)}(x)J_{m_1\times m_1}\end{pmatrix}$$

is the Schur complement [11] of $xI_{n_2}-A(G_2)$.

It is well known [5] that $R^TR=A(\mathcal{L}(G_1))+2I_{m_1}$, and that [5, Theorem 2.4.1] the eigenvalues of $\mathcal{L}(G_1)$ are $\lambda_i(G_1)+r_1-2$, for $i=1,2,\ldots,n_1$, and -2 repeated m_1-n_1 times, where $\mathcal{L}(G)$ denotes the line graph of G. Thus, by applying these arguments, the result follows from

$$\begin{aligned}\det(S) &= x^{n_1}\cdot\det\left(xI_{m_1}-\Gamma_{A(G_2)}(x)J_{m_1\times m_1}-\frac{1}{x}R^TR\right)\\ &= x^{n_1}\cdot\left\{\det\left(xI_{m_1}-\frac{1}{x}R^TR\right)-\Gamma_{A(G_2)}(x)\cdot\mathbf{1}_{m_1}^T\operatorname{adj}\left(xI_{m_1}-\frac{1}{x}R^TR\right)\mathbf{1}_{m_1}\right\}\\ &= x^{n_1}\cdot\left(1-\Gamma_{A(G_2)}(x)\cdot\mathbf{1}_{m_1}^T\left(xI_{m_1}-\frac{1}{x}R^TR\right)^{-1}\mathbf{1}_{m_1}\right)\cdot\det\left(xI_{m_1}-\frac{1}{x}R^TR\right)\\ &= x^{n_1}\cdot\Big(1-\Gamma_{A(G_2)}(x)\cdot\Gamma_{\frac{1}{x}R^TR}(x)\Big)\cdot\prod_{i=1}^{m_1}\left(x-\frac{2}{x}-\frac{1}{x}\lambda_i(\mathcal{L}(G_1))\right)\\ &= x^{m_1-n_1}\cdot\Big(x^2-2r_1-m_1x\cdot\Gamma_{A(G_2)}(x)\Big)\cdot\prod_{i=2}^{n_1}\Big(x^2-r_1-\lambda_i(G_1)\Big).\end{aligned}$$

□

Similar to Corollaries 2.2, 2.3 and 2.4, Theorem 3.1 implies the following results.

Corollary 3.2. [8, Theorem 1.2] *Let G_1 be an r_1-regular graphs on n_1 vertices and m_1 edges, and G_2 an r_2-regular graphs on n_2 vertices. Then the A-spectrum of $G_1 \underline{\vee} G_2$ consists of:*

(a) $\lambda_i(G_2)$, *for* $i = 2, 3, \ldots, n_2$;

(b) 0, *repeated* $m_1 - n_1$;

(c) $\pm\sqrt{r_1 + \lambda_j(G_1)}$, *for* $j = 2, 3, \ldots, n_1$;

(d) *the three roots of the equation*

$$x^3 - r_2x^2 - (m_1n_2 + 2r_1)x + 2r_1r_2 = 0.$$

Corollary 3.3. *Let G be an r-regular graph on n vertices and m edges with $m \geq n$, and let $p, q \geq 1$ be integers. Then the A-spectrum of $G \underline{\vee} K_{p,q}$ consists of:*

(a) 0, *repeated* $m - n + p + q - 2$ *times;*

(b) $\pm\sqrt{r + \lambda_i(G)}$, *for* $i = 2, 3, \ldots, n$;

(c) *the four roots of the equation*

$$x^4 - \big(pq + m(p + q) + 2r\big)x^2 - 2mpqx + 2pqr = 0.$$

Corollary 3.4. (a) *If G_1 and G_2 are A-cospectral regular graphs, and H is any graph, then $G_1 \underline{\vee} H$ and $G_2 \underline{\vee} H$ are A-cospectral.*

(b) *If G is a regular graph, and H_1 and H_2 are A-cospectral graphs with $\Gamma_{A(H_1)}(x) = \Gamma_{A(H_2)}(x)$, then $G \underline{\vee} H_1$ and $G \underline{\vee} H_2$ are A-cospectral.*

3.2 L-spectra of subdivision-edge joins

Theorem 3.5. *Let G_1 be an r_1-regular graph on n_1 vertices and m_1 edges, and G_2 an arbitrary graph on n_2 vertices. Then*

$$\begin{aligned}\phi\left(L(G_1 \underline{\vee} G_2); x\right) &= x \cdot (x - 2 - n_2)^{m_1 - n_1} \cdot \Big(x^2 - (2 + r_1 + m_1 + n_2)x + r_1n_2 + r_1m_1 + 2m_1\Big) \\ &\quad \cdot \prod_{i=2}^{n_2}\Big(x - m_1 - \mu_i(G_2)\Big) \cdot \prod_{i=2}^{n_1}\Big(x^2 - (2 + r_1 + n_2)x + r_1n_2 + \mu_i(G_1)\Big).\end{aligned}$$

Proof. Let R be the incidence matrix of G_1. Then the Laplacian matrix of $G_1 \underline{\vee} G_2$ can be written as

$$L(G_1 \underline{\vee} G_2) = \begin{bmatrix} r_1I_{n_1} & -R & 0_{n_1 \times n_2} \\ -R^T & (2 + n_2)I_{m_1} & -J_{m_1 \times n_2} \\ 0_{n_2 \times n_1} & -J_{n_2 \times m_1} & m_1I_{n_2} + L(G_2) \end{bmatrix}.$$

Thus the Laplacian characteristic polynomial of $G_1 \underline{\vee} G_2$ is given by

$$\begin{aligned}
\phi\left(L(G_1 \underline{\vee} G_2); x\right) &= \det \begin{bmatrix} (x-r_1)I_{n_1} & R & 0_{n_1 \times n_2} \\ R^T & (x-2-n_2)I_{m_1} & J_{m_1 \times n_2} \\ 0_{n_2 \times n_1} & J_{n_2 \times m_1} & (x-m_1)I_{n_2} - L(G_2) \end{bmatrix} \\
&= \det\big((x-m_1)I_{n_2} - L(G_2)\big) \cdot \det(S) \\
&= \det(S) \cdot \prod_{i=1}^{n_2} \Big(x - m_1 - \mu_i(G_2)\Big),
\end{aligned}$$

where

$$S = \begin{pmatrix} (x-r_1)I_{n_1} & R \\ R^T & (x-2-n_2)I_{m_1} - \Gamma_{L(G_2)}(x-m_1)J_{m_1 \times m_1} \end{pmatrix}$$

is the Schur complement [11] of $(x-m_1)I_{n_2} - L(G_2)$. Then, the result follows from

$$\begin{aligned}
\det(S) &= (x-r_1)^{n_1} \cdot \det\left((x-2-n_2)I_{m_1} - \Gamma_{L(G_2)}(x-m_1)J_{m_1 \times m_1} - \frac{1}{x-r_1}R^TR\right) \\
&= (x-r_1)^{n_1} \cdot \left\{\det\left((x-2-n_2)I_{m_1} - \frac{1}{x-r_1}R^TR\right)\right. \\
&\quad \left. -\Gamma_{L(G_2)}(x-m_1) \cdot \mathbf{1}_{m_1}^T \operatorname{adj}\left((x-2-n_2)I_{m_1} - \frac{1}{x-r_1}R^TR\right)\mathbf{1}_{m_1}\right\} \\
&= (x-r_1)^{n_1} \cdot \det\left((x-2-n_2)I_{m_1} - \frac{1}{x-r_1}R^TR\right) \\
&\quad \cdot \left(1 - \Gamma_{L(G_2)}(x-m_1) \cdot \mathbf{1}_{m_1}^T \left((x-2-n_2)I_{m_1} - \frac{1}{x-r_1}R^TR\right)^{-1} \mathbf{1}_{m_1}\right) \\
&= (x-r_1)^{n_1} \cdot \prod_{i=1}^{m_1}\left(x - 2 - n_2 - \frac{2}{x-r_1} - \frac{1}{x-r_1}\lambda_i(\mathcal{L}(G_1))\right) \\
&\quad \cdot \left(1 - \Gamma_{L(G_2)}(x-m_1)\Gamma_{\frac{1}{x-r_1}R^TR}(x-2-n_2)\right) \\
&= \frac{x}{x-m_1} \cdot (x-2-n_2)^{m_1-n_1} \cdot \prod_{i=2}^{n_1}\Big(x^2 - (2+r_1+n_2)x + r_1n_2 + \mu_i(G_1)\Big) \\
&\quad \cdot \Big(x^2 - (2+r_1+m_1+n_2)x + r_1n_2 + r_1m_1 + 2m_1\Big).
\end{aligned}$$

□

Theorem 3.5 implies the following results.

Corollary 3.6. (a) *If G_1 and G_2 are L-cospectral regular graphs, and H is an arbitrary graph, then $G_1 \underline{\vee} H$ and $G_2 \underline{\vee} H$ are L-cospectral.*

(b) *If G is a regular graph, and H_1 and H_2 are L-cospectral graphs, then $G \underline{\vee} H_1$ and $G \underline{\vee} H_2$ are L-cospectral.*

(c) *If* G_1 *and* G_2 *are* L*-cospectral regular graphs, and* H_1 *and* H_2 *are* L*-cospectral graphs, then* $G_1 \underline{\vee} H_1$ *and* $G_2 \underline{\vee} H_2$ *are* L*-cospectral.*

Corollary 3.7. *Let* G_1 *be an* r_1*-regular graph on* n_1 *vertices and* m_1 *edges, and* G_2 *an arbitrary graph on* n_2 *vertices. Then*

$$t(G_1 \underline{\vee} G_2) = \frac{(2+n_2)^{m_1-n_1} \cdot (r_1 n_2 + r_1 m_1 + 2m_1) \cdot \prod_{i=2}^{n_2} \Big(m_1 + \mu_i(G_2)\Big) \cdot \prod_{i=2}^{n_1} \Big(r_1 n_2 + \mu_i(G_1)\Big)}{m_1 + n_1 + n_2}.$$

3.3 Q-spectra of subdivision-edge joins

Theorem 3.8. *Let* G_1 *be an* r_1*-regular graph on* n_1 *vertices and* m_1 *edges, and* G_2 *an arbitrary graph on* n_2 *vertices. Then*

$$\begin{aligned} \phi\left(Q(G_1 \underline{\vee} G_2); x\right) &= (x-2-n_2)^{m_1-n_1} \cdot \prod_{i=1}^{n_2} \Big(x - m_1 - \nu_i(G_2)\Big) \\ &\quad \cdot \prod_{i=1}^{n_1-1} \Big(x^2 - (2+r_1+n_2)x + r_1 n_2 + 2r_1 - \nu_i(G_1)\Big) \\ &\quad \cdot \Big(x^2 - (2+r_1+n_2)x + r_1 n_2 - m_1(x-r_1) \cdot \Gamma_{Q(G_2)}(x-m_1)\Big). \end{aligned}$$

Proof. Let R be the incidence matrix of G_1. Then the signless Laplacian matrix of $G_1 \underline{\vee} G_2$ can be written as

$$Q(G_1 \underline{\vee} G_2) = \begin{bmatrix} r_1 I_{n_1} & R & 0_{n_1 \times n_2} \\ R^T & (2+n_2) I_{m_1} & J_{m_1 \times n_2} \\ 0_{n_2 \times n_1} & J_{n_2 \times m_1} & m_1 I_{n_2} + Q(G_2) \end{bmatrix}.$$

The rest of the proof is similar to that of Theorem 3.5 and hence we omit details. □

By applying (2.1) again, Theorem 3.8 implies the following result.

Corollary 3.9. *Let* G_1 *be an* r_1*-regular graphs on* n_1 *vertices and* m_1 *edges, and* G_2 *an* r_2*-regular graphs on* n_2 *vertices. Then*

$$\begin{aligned} \phi\left(Q(G_1 \dot{\vee} G_2); x\right) &= (x-2-n_2)^{m_1-n_1} \cdot \Big(x^3 - ax^2 + bx - 2r_1 r_2 n_2\Big) \cdot \prod_{i=1}^{n_2-1} \Big(x - m_1 - \nu_i(G_2)\Big) \\ &\quad \cdot \prod_{i=1}^{n_1-1} \Big(x^2 - (2+r_1+n_2)x + r_1 n_2 + 2r_1 - \nu_i(G_1)\Big). \end{aligned}$$

where $a = 2 + 2r_2 + r_1 + m_1 + n_2$ *and* $b = r_1 n_2 + 2m_1 + r_1 m_1 + 4r_2 + 2r_1 r_2 + 2r_2 n_2$*.*

Finally, Theorem 3.8 enables us to construct infinite Q-cospectral graphs.

Corollary 3.10. (a) *If* G_1 *and* G_2 *are* Q*-cospectral regular graphs, and* H *is a regular graph, then* $G_1 \underline{\vee} H$ *and* $G_2 \underline{\vee} H$ *are* Q*-cospectral.*

(b) *If* G *is a regular graph, and* H_1 *and* H_2 *are* Q*-cospectral graphs with* $\Gamma_{Q(H_1)}(x) = \Gamma_{Q(H_2)}(x)$*, then* $G \underline{\vee} H_1$ *and* $G \underline{\vee} H_2$ *are* Q*-cospectral.*

References

[1] S. Barik, S. Pati, B. K. Sarma, The spectrum of the corona of two graphs, SIAM J. Discrete Math. 24 (2007) 47–56.

[2] A. E. Brouwer, W. H. Haemers, Spectra of Graphs, Springer, 2012.

[3] S.-Y. Cui, G.-X. Tian, The spectrum and the signless Laplacian spectrum of coronae, Linear Algebra Appl. 437 (2012) 1692–1703.

[4] D. M. Cvetković, M. Doob, H. Sachs, Spectra of Graphs - Theory and Applications, Third edition, Johann Ambrosius Barth. Heidelberg, 1995.

[5] D. M. Cvetković, P. Rowlinson, H. Simić, An Introduction to the Theory of Graph Spectra, Cambridge University Press, Cambridge, 2010.

[6] I. Gopalapillai, The spectrum of neighborhood corona of graphs, Kragujevac J. Math. 35 (2011) 493–500.

[7] Y.-P. Hou, W.-C. Shiu, The spectrum of the edge corona of two graphs, Electron. J. Linear Algebra. 20 (2010) 586–594.

[8] G. Indulal, Spectrum of two new joins of graphs and infinite families of integral graphs, Kragujevac J. Math. 36 (2012) 133–139.

[9] C. McLeman, E. McNicholas, Spectra of coronae, Linear Algebra Appl. 435 (2011) 998–1007.

[10] S.-L. Wang, B. Zhou, The signless Laplacian spectra of the corona and edge corona of two graphs, Linear Multilinear Algebra (2012) 1–8, iFirst.

[11] F.-Z. Zhang, The Schur Complement and Its Applications, Springer, 2005.